Future of SR

Tetsuya Ishikawa RIKEN SPring-8 Center



Cheiron School



Plan

- This lecture, titled 'Future of SR', is designed to give you some background of SR history, then, discusses the possible future of SR facilities and sciences.
- Strategies for future may be different from person to person, or place to place, but important thing is to think about **YOUR** own future.
- Join us to make **NEW HISTORY OF SR** which is not the mere extrapolation from the present!

Brief Introduction of myself

- My name is Tetsuya Ishikawa. I am the Director of RIKEN SPring-8 Center and the Project Leader of X-ray Free Electron Laser.
- My research field is X-Ray Optics including those for coherent x-rays.
- I have been working around the SR facilities since 1982 when the Photon Factory started operation.
- I had been the Optics Group Leader/Beamline Division Director till 2006 when Dr. Shunji Goto took over the position.

You have learned a lot in this school ...



Thank you for joining us in this Cheiron School 2015. Now, you are attending the lecture titled "**Future of SR**".

You are learning a lot about synchrotron radiation and its applications in the preceding and following lectures. It is not surprising that you may feel the science and technologies related to the synchrotron radiation are too widely diverse to be an expertise in everything.

Don't be afraid, because no one can cover everything! You can be the world's top scientist in a certain field.

You have to know where you are...



All lectures given in this school are the maps of science/technology fields that have been already explored by other scientists. All of you, who are young with full potential, may want to construct new roads in the fields. So focus yourselves where you are on the map to start with.

New possibilities really lie in what has not been directed in the lectures: It is YOU to explore the new roads and to make the new SR history!

History can tell you something...



SR history has been a pursuit of Brilliance.

Short History of Synchrotron Radiation

1945 First (indirect) observation of SR; J. Blewett, G.E. 100 MeV betatron 1947 1st visual observation; G.E. 70 MeV synchrotron

ZEROth GENERATION SOURCES 1950's-60's: ELECTRON SYNCHROTRONS (cyclic accelerators)

<u>FIRST GENERATION SOURCES (storage rings)</u> 1970's: e+/e- COLLIDERS (Mostly Parasitic on High Energy Physics programs)

<u>SECOND GENERATION SOURCES</u> 1980's: NEW RINGS and FULLY DEDICATED USE OF e+/e- COLLIDERS, USE OF WIGGLERS & UNDULATORS

<u>THIRD GENERATION SOURCES</u> 1990's: LOW EMITTANCE RINGS WITH MANY STRAIGHT SECTIONS

<u>FOURTH GENERATION SOURCES</u> 2000's: LINAC-BASED SOURCES •Free-electron laser (FEL) •Energy Recovery Linac (ERL) DIFFRACTION-LIMITED RINGS; ULTRA-SHORT BUNCHES; NEW IDEAS

SR brightness has grown faster than the computing Power

Growth in X-Ray Brightness compared to growth in computing power



LBL/SSRL 30 Period Permanent Magnet Undulator -1980



In-Vacuum Permanent Magnet Undulator in SPring-8



Short-period in-vacuum undulator



$$\gamma = 2,000 \times E(GeV)$$

For $\lambda_{\text{photon}} = 0.1 \text{ nm}$: (*K*=2) $\lambda_{\text{magnet}} = 30 \text{ mm}, E = 11 \text{ GeV}$ $\lambda_{\text{magnet}} = 18 \text{ mm}, E = 8 \text{ GeV}$ cf. $\lambda_{\text{magnet}} = 100 \text{ um}, E = 600 \text{ MeV} !!$



Dr. Kitamura



In-vacuum undulator:

- Short-period undulator requires higher magnetic field for deflecting electrons.
 Magnets are located "in vacuum" for generating higher field with smaller gap (~ 3 mm).
- Variable gap undulator

In-vacuum undulators

- In-vacuum undulator technology was a turning point of the SR history.
- This is the reason why SPring-8 had been the world-largest synchrotron radiation facility for ~10 years.
- With the in-vacuum undulator technology, lower energy storage rings can emit x-rays from undulators.

Down-Sized SR Facilities

It's our tradition to make everything compact.



So is XFEL...

"X-ray Free Electron Laser, XFEL" coherent light to explore nano-world



Linac-Based Free Electron Laser Self-Amplified Spontaneous Emission (SASE)



Precedent XFEL Projects in the U.S. and EU

<u>US</u> SLAC: Stanford Linear Accelerator Center

- **Liniac Coherent Light Source : LCLS**
- Use Existing 2 mile Liniac
- Size: 2 km long
- Shortest Operating Wavelength: 0.15 nm
- To Start Operation in 2009
- RF gun/Normal Conducting Liniac/Out-of-Vacuum Undulator
- 3rd Priority in DOE's Future Facility Plan

<u>(EU)</u> DESY: Deutsches Elektronen-Synchrotron

European X-Ray Free-Electron Laser

- Collaboration among 12 EU Countries + China
- Size: 3.4 km long
- Shortest Operating Wavelength: 0.085 nm
- To Start Operation in 2013
- RF gun/Superconducting Liniac/Out-of-Vacuum Undulator



European XFEL

Japan's XFEL: SPring-8 Compact SASE Source (SCSS) Concept



8 GeV X-Ray Free Electron Laser Facility at SPring-8



A compact free-electron laser for generating coherent radiation in the extreme ultraviolet region

A list of authors and their affiliations appears at the end of the paper

Published online: 27 July 2008; doi:10.1038/nphoton.2008.134

Single-pass free-electron lasers based on self-amplified spontaneous emission¹⁻⁴ are enabling the generation of laser light at ever shorter wavelengths, including extreme ultraviolet⁵, soft X-rays and even hard X-rays⁶⁻⁸. A typical X-ray free-electron laser is a few kilometres in length and requires an electron-beam energy higher than 10 GeV (refs 6, 8). If such light sources are to become accessible to more researchers, a significant reduction in scale is desirable Here, we report observations of brilliant extreme-ultraviolet radiation from a 55-m-long compact self-amplified spontaneous-emission source, which combines short-period undulators with a highquality electron <u>source operating at a low acceleration energy of</u>



250 MeV. wavelengt energy of electron degraded dramatici achievem electron I Synch with wav

A down-sized design

Necessity is the mother of invention. Lasing in the extreme UV from a prototype compact free-electron-laser design is reported, continuing the push towards X-ray wavelengths.

Brian McNeil

is in the Department of Physics, University of Strathclyde, Glasgow G4 ONG, UK. e-mail: b.w.j.mcneil@strath.ac.uk

igh-gain free-electron lasers (FELs)

to observe phenomena at these scales also offers the enticing prospect of being able to control them. Recent progress towards these dreams has been made on a prototype of a Japanese X-ray FEL (XFEL) — the SPring-8 commissioning pencilled in for late 2013. Although the existing linear accelerator (Linac) at SLAC provided the LCLS team with a flying start, they have had the tricky task of adapting a system built at the end of the 1960s. Meanwhile,

My Personal View; Learning Old, Getting New



Industrial Applications

Some Tips

- Small or Big?
- Far or Near?
- One of you tools or Central Subject?
- Use your own brain.
- Find what has been not discussed, in reading papers or in listening to lectures.

Summary & Outlook

The future directions are not what you will be taught, but what you yourself shall find or create.

Thank you again for joining us this Cheiron School! We are looking forward to seeing you at some SR facility in near future.